

## Resource 1a

**Test Your Pregnancy Knowledge**

TRUE FALSE

- |  |                          |                          |
|--|--------------------------|--------------------------|
| 1. One out of five pregnancies begins in the very first month after a girl starts having sex.  | <input type="checkbox"/> | <input type="checkbox"/> |
| 2. When a male ejaculates into a female's vagina, about 50 million sperm are released.   | <input type="checkbox"/> | <input type="checkbox"/> |
| 3. Teen pregnancy is mostly a problem among minority girls.  | <input type="checkbox"/> | <input type="checkbox"/> |
| 4. There are only three days a month that a female can get pregnant. If you avoid having intercourse on those days, you are pretty safe.                           | <input type="checkbox"/> | <input type="checkbox"/> |
| 5. After a male ejaculates in the vagina, it takes about three minutes for the sperm to reach the uterus.  | <input type="checkbox"/> | <input type="checkbox"/> |
| 6. Sperm lives about six hours.  | <input type="checkbox"/> | <input type="checkbox"/> |
| 7. If a male uses the "withdrawal" method of birth control, the female won't get pregnant.   | <input type="checkbox"/> | <input type="checkbox"/> |
| 8. Half of teen pregnancies occur within the first six months after a girl begins to have intercourse.   | <input type="checkbox"/> | <input type="checkbox"/> |
| 9. You can't get pregnant if you've only engaged in heavy petting, but not intercourse.  | <input type="checkbox"/> | <input type="checkbox"/> |
| 10. If a female doesn't have an orgasm, she can't get pregnant.  | <input type="checkbox"/> | <input type="checkbox"/> |
| 11. You can't get pregnant during your period.   | <input type="checkbox"/> | <input type="checkbox"/> |
| 12. If a female forgets to take her birth control pill for a couple of days, she can take two the next day and be safe.  | <input type="checkbox"/> | <input type="checkbox"/> |
| 13. If a couple doesn't use birth control, they have a 50 percent chance of getting pregnant within one year.  | <input type="checkbox"/> | <input type="checkbox"/> |
| 14. Depo-provera is a shot taken every three months that offers good protection against pregnancy and has few side effects.  | <input type="checkbox"/> | <input type="checkbox"/> |
| 15. Emergency contraception, often called the "morning after pill," which is sometimes used by victims of rape, can be taken up to ten days after unprotected sex. | <input type="checkbox"/> | <input type="checkbox"/> |

Resource 1a, continued

TRUE FALSE

16. The pill can be rendered less or non-effective by certain antibiotics.
17. The NuvaRing is a contraceptive for guys which is put on the penis.
18. The Ortho-Evra patch is like a birth control pill, but absorbed through the skin.
19. Birth control devices like the NuvaRing or the diaphragm, offer significant STD protection as well as pregnancy prevention.
20. The IUD is a long lasting intrauterine device that a health provider inserts through the cervix into the uterus.